


REPORT DOCUMENTATION PAGE			<i>Form Approved</i> OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.				
1. REPORT DATE (DD-MM-YYYY) 2012		2. REPORT TYPE Open Literature		3. DATES COVERED (From - To)
4. TITLE AND SUBTITLE Reversal of BoNT/A-mediated inhibition of muscle paralysis by 3,4-diaminopyridine and roscovitine in mouse phrenic nerve-hemidiaphragm preparations		5a. CONTRACT NUMBER		
		5b. GRANT NUMBER		
		5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Adler, M., Deshpande, S.S., Apland, J.P., Murray, B., Borrell, A.		5d. PROJECT NUMBER		
		5e. TASK NUMBER		
		5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: MCMR-CDT-N 3100 Ricketts Point Road		8. PERFORMING ORGANIZATION REPORT NUMBER USAMRICD-P12-011		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Defense Threat Reduction Agency 8725 John J. Kingman Road STOP 6201 Fort Belvoir, VA 22060-6201		10. SPONSOR/MONITOR'S ACRONYM(S)		
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				
13. SUPPLEMENTARY NOTES Published in Neurochemistry International, 61(6), 866-873, 2012. This research was supported by the Defense Threat Reduction Agency- Joint Science and Technology Office, Medical S&T Division.				
14. ABSTRACT See reprint.				
15. SUBJECT TERMS Botulinum neurotoxin, 3,4-Diaminopyridine, Roscovitine, Acetylcholine, Mouse, Diaphragm muscle, Aminopyridine, Potassium channel blocker, Calcium channel agonist, Bioterrorism				
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UNLIMITED	18. NUMBER OF PAGES 8
a. REPORT UNCLASSIFIED	b. ABSTRACT UNCLASSIFIED	c. THIS PAGE UNCLASSIFIED		
				19a. NAME OF RESPONSIBLE PERSON Michael Adler
				19b. TELEPHONE NUMBER (include area code) 41-436-1913



Reversal of BoNT/A-mediated inhibition of muscle paralysis by 3,4-diaminopyridine and roscovitine in mouse phrenic nerve-hemidiaphragm preparations [☆]

Michael Adler ^{a,b,*}, Sharad S. Deshpande ^a, James P. Aplan ^a, Bridget Murray ^a, Andrew Borrell ^a

^a Neurobehavioral Toxicology Branch, Analytical Toxicology Division, U.S. Army Medical Research Institute of Chemical Defense, 3100 Ricketts Point Road, APG, MD 21010-5400, USA

^b Department of Chemistry and Biochemistry, University of Massachusetts, 285 Old Westport Road, Dartmouth, MA 02747-2300, USA

ARTICLE INFO

Article history:

Available online 25 July 2012

Keywords:

Botulinum neurotoxin
3,4-Diaminopyridine
Roscovitine
Acetylcholine
Mouse
Diaphragm muscle
Aminopyridine
Potassium channel blocker
Calcium channel agonist
Bioterrorism

ABSTRACT

Botulinum neurotoxins (BoNTs) comprise a family of neurotoxic proteins synthesized by anaerobic bacteria of the genus *Clostridium*. Each neurotoxin consists of two polypeptide chains: a 100 kDa heavy chain, responsible for binding and internalization into the nerve terminal of cholinergic motoneurons and a 50 kDa light chain that mediates cleavage of specific synaptic proteins in the host nerve terminal. Exposure to BoNT leads to cessation of voltage- and Ca²⁺-dependent acetylcholine (ACh) release, resulting in flaccid paralysis which may be protracted and potentially fatal.

There are no approved therapies for BoNT intoxication once symptoms appear, and specific inhibitors of the light chain developed to date have not been able to reverse the consequences of BoNT intoxication. An alternative approach for treatment of botulism is to focus on compounds that act by enhancing ACh release. To this end, we examined the action of the K⁺ channel blocker 3,4-diaminopyridine (3,4-DAP) in isolated mouse hemidiaphragm muscles intoxicated with 5 pM BoNT/A. 3,4-DAP restored tension within 1–3 min of application, and was effective even in totally paralyzed muscle. The Ca²⁺ channel activator (R)-roscovitine (Ros) potentiated the action of 3,4-DAP, allowing for use of lower concentrations of the K⁺ channel blocker. In the absence of 3,4-DAP, Ros was unable to augment tension in BoNT/A-intoxicated muscle. This is the first report demonstrating the efficacy of the combination of 3,4-DAP and Ros for the potential treatment of BoNT/A-mediated muscle paralysis.

Published by Elsevier Ltd.

1. Introduction

The seven serotypes of botulinum neurotoxin (BoNT) are the most potent substances in nature, and exposure to as little as 1–3 ng/kg may be sufficient to cause human lethality (Arnon et al., 2001; Lindström and Korkeala, 2006; Rega et al., 2010). The neurotoxins are produced by spore forming anaerobic bacteria, chiefly *Clostridium botulinum*, and by a limited number of other clostridial strains (Simpson, 2004; Sobel, 2005). BoNTs are secreted initially as relatively inactive ~150 kDa protoxins (range 140–167 kDa),

[☆] The views expressed are those of the authors and do not reflect the official policy of the U.S. Department of Army, Department of Defense, or U.S. Government. The experimental protocol was approved by the animal Care and Use Committee at the United States Army Medical Research Institute of Chemical Defense, and all procedures were conducted in accordance with the principles stated in the Guide for the Care and Use of Laboratory Animals and the Animal Welfare Act of 1966 (P.L. 89-544), as amended. This research was supported by the Defense Threat Reduction Agency-Joint Science and Technology Office, Medical S & T Division.

* Corresponding author at: Neurobehavioral Toxicology Branch, Analytical Toxicology Division, U.S. Army Medical Research Institute of Chemical Defense, 3100 Ricketts Point Road, APG, MD 21010-5400, USA. Tel.: +1 410 436 1913; fax: +1 410 436 2750.

E-mail addresses: michael.adler2@us.army.mil, miklos78@hotmail.com (M. Adler).

surrounded by a complex of neurotoxin-associated proteins that protect BoNT from degradation in the gastrointestinal tract (Kukreja and Singh, 2007; Gu et al., 2012). The protoxin is subsequently cleaved to form the active dichain neurotoxin, consisting of a 100 kDa heavy chain (HC) and a 50 kDa light chain (LC) (DasGupta and Sugiyama, 1972).

The BoNTs have three functional domains: binding, translocation and catalytic. Binding is mediated by the C-terminal region of the heavy chain, which interacts with gangliosides and protein receptors located on cholinergic nerve terminals (Dong et al., 2006; Benson et al., 2011); selective binding of BoNT to these receptors is responsible for the cholinergic selectivity of the BoNTs. The N-terminal region of the HC promotes translocation of the LC into the cytosol (Korazova and Montal, 2003; Fischer and Montal, 2007). The LC is a Zn²⁺-containing endopeptidase that cleaves specific sites on SNARE (soluble-N-ethylmaleimide sensitive factor attachment protein receptor) proteins, leading to a cessation of evoked transmitter release (Montecucco et al., 2005).

All seven serotypes of BoNT (A–G) inhibit release of acetylcholine (ACh) from cholinergic nerve terminals that innervate skeletal muscle, autonomic ganglia and post-ganglionic parasympathetic organs (Simpson, 2004). In skeletal muscle, inhibition of transmitter release leads to flaccid paralysis, which can progress to generalized

muscle weakness and death when the muscles of respiration become sufficiently compromised.

Early signs of BoNT intoxication (botulism) include visual disturbances, difficulties in swallowing and impairment of speech (Sobel, 2005). At this stage, botulism can be treated by infusion of serotype specific antitoxin. However, when symptoms progress to generalized paralysis, antitoxin is no longer effective, and patients will need treatment in an intensive care facility (Tacket et al., 1984; Hatheway et al., 1984). The more severe cases may also require extensive periods of artificial ventilation, enteral feeding and physical therapy after discharge from intensive care (Shapiro et al., 1998; Robinson and Nahata, 2003; Marcus, 2009).

BoNT has numerous attributes that make it appealing to terrorists, including an unusually high potency, long duration of action and the potential to cause widespread panic with major social disruption (Arnon et al., 2001; Robinson and Nahata, 2003). Accordingly, BoNT has been classified as a Tier 1 select toxin by the U.S. Department of Health and Human Services, the only non-infectious agent to receive this designation.

Of the seven serotypes, BoNT/A has the highest potency and longest duration of action and it therefore represents the greatest bioterrorist threat (Arnon et al., 2001). A number of complementary approaches have been used to develop pharmacological antagonists for BoNT/A: these consist of synthesis and screening of small molecule inhibitors (SMLs) for their ability to inactivate the BoNT/A LC, strategies to enhance degradation of the LC from intoxicated nerve terminals and evaluation of physiological antagonists such as the K^+ channel blocker 3,4-diaminopyridine (3,4-DAP) and the Ca^{2+} channel activator (R)-roscovitine (Ros) that can overcome the inhibitory action of BoNT and restore ACh release. To date, SMLs have only been able to slow the rate of BoNT/A-mediated paralysis, but they have not proved effective in reversing the paralytic action of BoNT, and strategies to accelerate the removal of BoNT/A LC from the nerve terminal have not progressed beyond the proof-of-concept stage (Tsai et al., 2010). However, physiological antagonists have shown remarkable promise both in slowing the onset of muscle paralysis and in restoring tension in BoNT/A-paralyzed muscles (Molgo et al., 1980, 1987; Adler et al., 1995, 1996; Mayorov et al., 2010).

A notable advantage of developing 3,4-DAP and Ros for treatment of botulism is that both have an extensive history of clinical use: aminopyridines for the symptomatic treatment of multiple sclerosis, Lambert Eaton Myasthenic Syndrome (LEMS) and downbeat nystagmus (Shi and Sun, 2011; Kalla et al., 2011), and Ros for breast, ovarian and prostate cancer (Benson et al., 2007; Yarotsky and Elmslie, 2012). In addition, 3,4-DAP and other K^+ channel blockers have been investigated for treatment of human botulism, with some success (Puggiari and Cherington, 1978; Kalia and Swartz, 2011).




Fig. 1. Structure of the K^+ channel blocker 3,4-DAP and the Ca^{2+} channel activator Ros. 3,4-DAP (CAS No. 54-96-6) is a *N*-heterocyclic tertiary amine also known as amifampridine in clinical use. The phosphate salt was licensed in Europe as an orphan drug for treatment of rare muscle disorders such as LEMS in 2010. Ros (CAS No. 186692-46-6) is a 2,6,9-tri substituted purine analog that is marketed under the proprietary name Seliciclib primarily as an antitumor agent.

The current study was undertaken to examine the ability of 3,4-DAP and Ros (Fig. 1) to reverse muscle paralysis produced by BoNT/A in isolated mouse hemidiaphragm muscle. The results indicate that 3,4-DAP can restore muscle tension completely, and combinations of 3,4-DAP with Ros allow for use of lower, less toxic concentrations of the former. It is concluded that 3,4-DAP and Ros are promising lead compounds for the development of medical countermeasures for botulism. This is the first study examining the efficacy of Ros in BoNT intoxication, as well as the first showing efficacy of co-administering Ros and 3,4-DAP in reversing BoNT/A-mediated muscle paralysis.

2. Methods

2.1. Muscle preparation

Experiments were performed *in vitro* on isolated hemidiaphragm muscles dissected from adult male CD-1 mice (19–24 g on arrival; Charles River Laboratories, Wilmington, MA, USA). Mice were housed in facilities approved by the Association for Assessment and Accreditation of Laboratory Animal Care International (AAALAC) with food and water provided *ad libitum*. Animals were euthanized by decapitation following exposure to excess isoflurane. Hemidiaphragms with attached phrenic nerves were mounted in 20-ml tissue baths containing Tyrode's solution of the following composition (mM): NaCl, 137; KCl, 5; $MgSO_4$, 1; $NaHCO_3$, 24; NaH_2PO_4 , 1; $CaCl_2$, 1.8, and glucose, 11. The solution was bubbled with a gas mixture of 95% O_2 /5% CO_2 yielding a pH of 7.3–7.4. Resting tension was maintained at 0.7 g to generate optimal nerve-evoked contractions.

2.2. Tension recordings

Twitch tension was elicited by supramaximal stimulation of the phrenic nerve via bipolar, stainless steel electrodes (6.0–9.0 V, 0.2 ms duration) at 0.033 Hz. Tetani were elicited by repetitive stimulation at 30 Hz for 1 s, with 1-min intervals between stimulus trains. Muscle tensions were measured using Grass FT03 force displacement transducers (West Warwick, RI, USA), digitized and analyzed offline using pClamp software v. 10.1 (Molecular Devices, Sunnyvale, CA, USA). Following a 15- to 20-min equilibration, muscles were exposed to 5 pM of pure BoNT/A (MetabioLogics, Inc., Madison, WI, USA) for 30 min at room temperature (18–22 °C) in the absence of nerve stimulation. The muscles were then washed with control Tyrode's solution to remove unbound BoNT/A, warmed to 36 °C and monitored for development of paralysis. 3,4-DAP (Sigma–Aldrich, St. Louis, MO, USA) and Ros (A.G. Scientific, Inc., San Diego, CA, USA) were generally added to the bath when muscle tensions declined to ~50% of their initial values. Tensions were recorded for 1–3 h after drug addition to determine the time course for reversal of muscle paralysis.

2.3. Drug preparation

Stock solutions of 3,4-DAP were prepared in deionized water at concentrations of 20 mM. Stock solutions of Ros were prepared in dimethylsulfoxide (DMSO) at a concentration of 50–100 mM and stored in opaque vials to limit their photoreactivity. Stock solutions were stored at 4 °C.

2.4. Data analysis

Unless stated otherwise, all data are expressed as means \pm SEM. Statistical analysis was performed using a one-way analysis of variance (ANOVA) followed by the Bonferroni multiple comparison




Fig. 2. Recordings of neurally-elicited twitch tensions in mouse hemidiaphragm muscle. (A) The muscle was intoxicated with BoNT/A (5 pM) at 0-time (short arrow), and 3,4-DAP (10 μ M) was added when tension had declined to 17% of control. 3,4-DAP (10 μ M) produced a partial restoration of tension. Tensions were normalized by averaging 10 control responses just prior to BoNT exposure, and setting these equal to 1. (B) Recordings illustrating the effects of Ros on a hemidiaphragm muscle intoxicated with 5 pM BoNT/A (short arrow). Ros was added in increasing concentrations at the indicated time points. Addition of 10 μ M 3,4-DAP in the presence of 100 μ M Ros led to complete restoration of tension. Tensions were normalized by averaging 10 control responses just prior to BoNT/A, and setting these equal to 1. Contractions elicited 2.5–40 min after addition of BoNT/A were omitted for clarity.

test (GraphPad, InStat, La Jolla, CA, USA). $P < 0.05$ was considered to be statistically significant.

3. Results

3.1. Time course of BoNT/A-mediated paralysis and reversal by 3,4-DAP

Twitch tension in control mouse hemidiaphragm muscle ranged from 0.7 to 2.2 g and was stable with time, decreasing by less than 10% over 4 h. Exposure to 5 pM BoNT/A led to a progressive reduction in tension after a variable latent period, culminating in total muscle paralysis (half-time = 61.7 ± 5.8 min) (Fig. 2). In this hemidiaphragm, 10 μ M 3,4-DAP was added 77 min after 5 pM BoNT/A, when twitch tension had declined to less than 20% of control. Tension increased within 2 min of 3,4-DAP addition and continued to rise for the next 15 min before attaining steady-state at 45% of control. The ability of 3,4-DAP to reverse muscle paralysis is of considerable interest since no SMI examined to date has been able to increase tension in BoNT/A-intoxicated muscle after onset of paralysis (Adler and Nicholson, 2008; Li et al., 2010; Ruthel et al., 2011).

3.2. Effect of Ros and Ros–3,4-DAP combination on muscle tension

In the absence of 3,4-DAP, Ros was unable to halt the progression of BoNT/A-mediated depression of twitch tension at any

concentration tested (10–100 μ M). Fig. 2B illustrates the effect of Ros in a hemidiaphragm muscle intoxicated with 5 pM BoNT/A. Addition of 10 or 30 μ M Ros led to a transient increase in tension, followed by resumption of paralysis. Raising the concentration of Ros to 100 μ M failed to produce even a transient increase, and twitches continued to decline to a level where they could barely be distinguished from baseline noise (Fig. 2B). A subsequent addition of 10 μ M 3,4-DAP to this preparation at a time of near complete muscle paralysis produced a rapid restoration of tension. From comparison of Fig. 2A and B, it is clear that Ros markedly augments the action of 10 μ M 3,4-DAP in BoNT/A-intoxicated muscle.

Data illustrating the interaction of 3,4-DAP and Ros in 11–16 hemidiaphragm muscles are shown in Fig. 3. In control muscle, twitch tension was potentiated after addition of either 30 μ M Ros or 10 μ M 3,4-DAP (Fig. 3A): the former to 128% of control ($P < 0.05$) and the latter to 230% of control ($P < 0.001$). Addition of 30 μ M Ros to the Tyrode's solution containing 10 μ M 3,4-DAP elevated tensions to 287% of control ($P < 0.05$, combination relative to 3,4-DAP alone).

The individual drugs and combination were subsequently tested in muscles intoxicated with 5 pM BoNT/A (Fig. 3B). The drugs were added after muscle tensions had declined to $54 \pm 4\%$ of control and evaluated for their ability to restore tension after 20- to 30-min of incubation. No increase in twitch tension was observed in BoNT/A-intoxicated muscle in the presence of 30 μ M Ros (Fig. 3B), consistent with the data in Fig. 2B. In contrast, addition of




Fig. 3. Histograms showing the effects of 30 μ M Ros, 10 μ M 3,4-DAP and combination of the two drugs in control hemidiaphragm muscles (A) and in hemidiaphragm muscles intoxicated with 5 pM BoNT/A (B). Data shown are mean \pm SEM of supramaximal twitch tensions obtained from 11 to 16 muscles per condition. Recording were made 20–30 min after drug addition. Muscles were washed for 40 min with control Tyrode's solution between Ros and 3,4-DAP addition. (A) Differs significantly from control (* P < 0.05, $^{\dagger}P$ < 0.001); differs significantly from values observed in 10 μ M 3,4-DAP ($^{\dagger}P$ < 0.05). (B) Differs significantly from control (* P < 0.001); differs significantly from values in the presence of BoNT/A (** P < 0.05, $^{\#}P$ < 0.001); differs significantly from values observed in 10 μ M 3,4-DAP ($^{\dagger}P$ < 0.001).

10 μ M 3,4-DAP increased muscle tension by 26% (P < 0.05). Addition of 30 μ M Ros to bathing medium containing 10 μ M 3,4-DAP led to an even greater increase in tension (P < 0.001, combination vs. 3,4-DAP alone). Co-application of these two drugs was thus able to restore tension to near control levels (Fig. 3B).

3.3. Effects of high concentrations of 3,4-DAP

Since 3,4-DAP can reverse the inhibitory action of BoNT/A on its own, it may be questioned as to whether addition of a second drug is necessary. Although 3,4-DAP is fully effective (Molgó et al., 1980), the high concentrations required generally result in abnormal contractions, as depicted in Fig. 4. The traces were obtained from a hemidiaphragm muscle under control conditions and after sequential additions of BoNT/A and 3,4-DAP at the indicated concentrations. Exposure of the muscle to 5 pM BoNT/A led to a profound reduction in twitch tension (Fig. 4B) as noted earlier (Fig. 2). Addition of 30 μ M 3,4-DAP caused a marked enhancement of tension with no change in the time course (Fig. 4C). Raising the 3,4-DAP concentration to 100 μ M led to a further increase in twitch tension, accompanied by abnormal repetitive after-discharges (Fig. 4D). Such after-discharges act to prolong the relaxation phase of twitches, thus interfering with the ability of muscles to follow higher stimulation frequencies. By combining 3,4-DAP with Ros, it was possible to reverse the paralytic effect of BoNT/A without altering the normal characteristics of muscle twitches (Fig. 2B).

3.4. Effect of 3,4-DAP and Ros on tensions elicited by repetitive stimulation

Since tension in skeletal muscle is normally generated by repetitive firing of motoneurons, it was of interest to evaluate the actions of 3,4-DAP, Ros and the combination of the two drugs under conditions of repetitive stimulation. Data obtained with 30 Hz trains are shown in Fig. 5. This frequency was selected because it is within the physiological range for mouse phrenic motoneurons (Paton, 1997). In response to 30 Hz stimulation, control muscles exhibited a sustained plateau, with superimposed cycles of contraction and relaxation (Fig. 5A). Exposure to 5 pM BoNT/A led to a reduction in tetanic tension amplitude and area (Fig. 5B). Addition of 30 μ M Ros to the BoNT/A-intoxicated muscle resulted in a small increase in tetanic tension (Fig. 5C), and a more pronounced increase was observed after addition of 10 μ M 3,4-DAP (Fig. 5D). As with twitch tensions, 30 μ M Ros potentiated the action

of 10 μ M 3,4-DAP (Fig. 5E). A high concentration of 3,4-DAP (100 μ M) elicited an even more dramatic enhancement of tetanic tension; however, both the plateau and baseline following the train exhibited instability (Fig. 5F).

Data from 5 to 9 such experiments are shown in Fig. 6. The ability of Ros and 3,4-DAP to reverse the action of 5 pM BoNT was assessed after tetanic area was reduced to $51 \pm 3\%$ of control. Addition of 30 μ M Ros increased tetanic area by a small but not significant amount (19%), whereas 10 μ M 3,4-DAP augmented tetanic area by 61% (P < 0.05). Addition of Ros to the 3,4-DAP-containing solution led to a further increase, such that in the presence of both drugs, tetanic area was $122 \pm 12\%$ of control (P < 0.01, combination vs. 3,4-DAP).

4. Discussion

The results of this investigation demonstrate that K^+ channel blockers and Ca^{2+} channel activators have important roles as potential medical countermeasures for the treatment of botulism. Of the two classes of drugs, the K^+ channel blockers have been studied more extensively. Their antagonism of BoNT/A-mediated paralysis stems from their ability to prolong the duration of the nerve terminal action potential (Penner and Dreyer, 1986), leading to a greater influx of Ca^{2+} during nerve stimulation. Coupling of increased Ca^{2+} influx to nerve impulses enables the K^+ channel blockers to produce striking increases in the amplitude of endplate potentials (EPPs) and of nerve-evoked twitch tensions (Lundh et al., 1977; Adler et al., 1979, 1995).

4.1. Role of K^+ channel blockers in antagonizing BoNT/A-mediated muscle paralysis

A number of K^+ channel blockers have been evaluated for their ability to antagonize the actions of BoNT, including guanidine, 4-aminopyridine, 3,4-DAP and tetraethylammonium (Cherington and Ryan, 1968; Cherington and Schultz, 1977; Lundh et al., 1977; Molgó et al., 1980, 1987; Simpson, 1986). Aminopyridines and tetraethylammonium inhibit different K^+ channel subtypes at the mammalian motor nerve terminal (Penner and Dreyer, 1986; Lin and Lin-Shiau, 1997; Nakamura and Takahashi, 2007), and both would appear to be useful for counteracting the inhibitory action of BoNT on transmitter release. Of the K^+ channel blockers examined, the most promising candidate thus far has been 3,4-DAP. 4-Aminopyridine produced a higher incidence of undesirable central




Fig. 4. Traces showing the effect of increasing concentrations of 3,4-DAP on the amplitude and time course of single twitches. The records were obtained from a hemidiaphragm muscle intoxicated with 5 pM BoNT/A and incubated sequentially with 30 and 100 μ M 3,4-DAP. The time course of twitches were unaltered by BoNT or 30 μ M 3,4-DAP, whereas 100 μ M 3,4-DAP produced a series of after-discharges that prolonged the relaxation phase.

nervous system (CNS) side effects, presumably because of its greater lipid solubility (Molgó et al., 1980). Tetraethylammonium, although devoid of CNS toxicity, has been found to cause a marked inhibition of muscle ACh receptor channels, thereby exacerbating BoNT/A-mediated muscle paralysis (Adler et al., 1979, 1995; Thesleff, 1989). When added to nerve-muscle preparations prior to BoNT, 3,4-DAP produced a marked delay in the time-to-block of nerve-evoked muscle contractions (Simpson, 1986; Mayorov et al., 2010). When applied after BoNT, 3,4-DAP was able to elevate tensions to control values or above (Figs. 2 and 3). In addition, unlike other putative BoNT/A antagonists, 3,4-DAP could restore tension even in muscles that were paralyzed for 2 weeks prior to drug administration (Adler et al., 1996).

4.2. Limitations of K^+ channel blockers in BoNT intoxication

In spite of these successes, 3,4-DAP also has limitations: (1) the efficacy of 3,4-DAP is largely confined to serotype A (Simpson, 1986; Adler et al., 1996), (2) concentrations of 3,4-DAP that are effective in reversing BoNT/A intoxication are toxic to humans (Bever et al., 1990; Vollmer et al., 2009), and (3) 3,4-DAP has a brief *in vivo* half-life relative to that of BoNT/A (Adler et al., 1996). The short duration of action can be offset, however, by use of an infusion delivery as demonstrated by Adler et al. (2000) with subcutaneously implanted osmotic minipumps. In addition, slow release formulations of the aminopyridines are currently available (Korenke et al., 2008; Vollmer et al., 2009; Hayes, 2011).

The basis for the lack of response to 3,4-DAP by the other serotypes is not well understood. At a functional level, BoNT/A-intoxicated neuromuscular junctions undergo an attenuated but synchronous release of ACh following stimulation; preparations intoxicated by serotypes B, D and F produce asynchronous release where the ACh quanta are dispersed and cannot summate to produce suprathreshold EPPs (Lundh et al., 1977; Molgó et al., 1980; Thesleff, 1989).

It is readily apparent that the lack of synchrony would prevent 3,4-DAP from restoring transmitter release; however, the factors that lead to asynchronous release are still not well understood. In this context, it is of interest that homozygous SNAP-25 null neurons were able to generate only asynchronous EPSCs; fast synchronous EPSCs could be restored, however, by lentiviral transfection with SNAP-25 (Delgado-Martínez et al., 2007). Collectively, these observations suggest that synchronous release requires SNAP-25, the substrate for BoNT/A LC, as well as intact synaptobrevin, the substrate for BoNT/B, /D and /F (Simpson, 2004).

4.3. K^+ channel blockers in human botulism

The K^+ channel blockers guanidine and 3,4-DAP have been evaluated in a small number of human botulism cases. The general findings were that while these blockers produced some increase in muscle strength, their use did not lead to the return of spontaneous ventilation (Cherington and Schultz, 1977; Davis et al., 1992). It is not clear if respiratory muscles are less responsive to K^+ blockers than limb or postural muscles or if the doses used clinically were insufficient to reverse muscle paralysis (Davis et al., 1992). High doses of 3,4-DAP were not attempted in patient studies to avoid the risk of seizures and other potential side effects. Based on the marked differences in 3,4-DAP concentrations that are effective in reversing BoNT/A intoxication in isolated diaphragm muscle ($\geq 10 \mu$ M; Figs. 2A and 3B), and the plasma levels in patients receiving the maximum tolerated dose of 3,4-DAP for conditions such as amyotrophic lateral sclerosis ($1.2 \pm 0.5 \mu$ M) (Aisen et al., 1995), it is likely that BoNT-intoxicated patients did not receive a dose that was sufficient to reverse paralysis in the case reports of Cherington and Schultz (1977) or Davis et al. (1992).

4.4. Development of more effective K^+ channel blockers

At the present time, the K^+ blockers hold promise as potential therapeutic agents, but additional strategies, such as development




Fig. 5. Effect of 3,4-DAP and Ros on tetanic tensions elicited with 1-s long trains at 30 Hz. Tensions (amplitude and area) were depressed after exposure to 5 pM BoNT/A (B). Tensions were increased slightly by 30 μ M Ros (C) but more substantially by 10 μ M 3,4-DAP (D). The combination of 30 μ M Ros and 10 μ M 3,4-DAP raised tensions to values exceeding control (E). Tetanic tensions were enhanced even more markedly by 100 μ M DAP; however, this concentration also produced tetanic fade and baseline instability (F).

of more selective compounds, and targeting of the inhibitors to neuromuscular and neuroeffector synapses, will be required to exploit their full potential. With regard to more selective inhibitors, Mayorov et al. (2010) synthesized new analogs of 3,4-DAP with the goal of finding compounds that displayed both an enhanced affinity for nerve terminal K^+ channels and a reduced propensity to cross the blood brain barrier. Although, none of the analogs were more potent than 3,4-DAP, one was found to have a more favorable peripheral to CNS distribution (Mayorov et al., 2010). Complicating the search for aminopyridines with low CNS toxicity is that their binding site on the K^+ channel is accessible only from the cytoplasmic membrane surface (Howe and Ritchie, 1991; Muñoz-Caro and Niño, 2002). This makes the goal of finding compounds with reduced CNS penetration challenging, since such compounds would also have an impaired ability to gain access to the cytoplasmic surface of the membrane. For this reason, it may be profitable to also consider K^+ channel blockers that act on the outer surface of the membrane in future studies.

4.5. Combination of K^+ channel blockers with Ca^{2+} channel activators

To address the issue that only high and potentially toxic doses of 3,4-DAP can antagonize the actions of BoNT, we examined the effect of combining 3,4-DAP with Ros, a trisubstituted purine analog best known for its action as a cyclin-dependent kinase (CDK) inhibitor (Fig. 1) (Benson et al., 2007). Distinct from its action on cell cycle regulation, Ros is also a novel activator of neuronal N-, P/Q- and R-type Ca^{2+} channels, and acts by slowing channel deactivation, thus prolonging channel open state (Buraei et al., 2005, 2007). These channels have been implicated in the regulation of transmitter release (Yan et al., 2002; DeStefino et al., 2010). Ros has been examined for treatment of human immunodeficiency virus type-1 (Guendel et al., 2010) and advanced malignancies (Le Tourneau et al., 2010), and is in Phase II trials for non-small cell lung cancer and nasopharyngeal carcinoma (Aldoss et al., 2009). Importantly, the plasma concentrations of Ros found in clinical trials are in the range of those used in the present study (Benson et al., 2007).




Fig. 6. Histograms showing the effects of 30 μ M Ros, 10 μ M 3,4-DAP and combination of the two drugs in hemidiaphragm muscles intoxicated with 5 pM BoNT/A. Data shown are mean \pm SEM of integrated tension records obtained from 5 to 9 hemidiaphragm muscles per condition. Recording were made 20–30 min after drug addition. Ros and 3,4-DAP were added to separate muscles, and combinations were achieved by subsequent addition of Ros or 3,4-DAP, as appropriate. Differs significantly from control (* P < 0.001); differs significantly from values in the presence of BoNT/A (** P < 0.05, # P < 0.001); differs significantly from values observed in 10 μ M 3,4-DAP († P < 0.01).

Unlike 3,4-DAP, Ros was not able to reverse BoNT-mediated depression of twitch tension on its own. However, when 30 μ M Ros and 10 μ M 3,4-DAP were co-applied, muscle tensions were restored rapidly to near control levels for twitch tension (Fig. 3) and above control for tetanic tension (Fig. 6). These results are encouraging since they demonstrate the possibility of achieving a prompt recovery from paralysis by using drugs with synergistic mechanisms of action: increased Ca^{2+} influx via K^+ channel blockade (Molgó et al., 1980) and enhanced Ca^{2+} conductance via prolongation of the channel open time (Yan et al., 2002; Buraei et al., 2005, 2007). Although the concentrations of 3,4-DAP used in this study (10–100 μ M) are toxic systemically (Aisen et al., 1995), the concentration of Ros falls within values for serum levels that are well-tolerated by patients (Benson et al., 2007). Based on the success of targeting Ca^{2+} entry by two complementary approaches, it is reasonable to assume that by making even modest improvements in the margin of safety of the K^+ channel blocker and in the potency of the Ca^{2+} channel activator, we should be able to develop a safe and effective treatment for BoNT/A intoxication.

Acknowledgements

The authors would like to thank Patricia Mangel and Fiona Brabazon for expert technical assistance and James D. Nicholson and Richard E. Sweeney for many helpful discussions.

References

- Adler, M., Capacio, B., Deshpande, S.S., 2000. Antagonism of botulinum toxin A-mediated muscle paralysis by 3,4-diaminopyridine delivered via osmotic minipumps. *Toxicon* 38, 1381–1388.
- Adler, M., MacDonald, D.A., Sellin, L.C., Parker, G.W., 1996. Effect of 3,4-diaminopyridine on rat extensor digitorum longus muscle paralyzed by local injection of botulinum neurotoxin. *Toxicon* 34, 237–249.
- Adler, M., Nicholson, J.D., 2008. Evaluation of toosendanin as a botulinum neurotoxin antagonist. *The Botulinum J.* 1, 208–218.
- Adler, M., Oliveira, A.C., Albuquerque, E.X., Mansour, N.A., Eldefrawi, A.T., 1979. Reaction of tetraethylammonium with the open and closed conformation of the acetylcholine receptor ionic channel complex. *J. Gen. Physiol.* 74, 129–152.
- Adler, M., Scovill, J., Parker, G., Lebeda, F.J., Piotrowski, J., Deshpande, S.S., 1995. Antagonism of botulinum toxin-induced muscle weakness by 3,4-diaminopyridine in rat phrenic nerve-hemidiaphragm preparations. *Toxicon* 33, 527–537.
- Aisen, M.L., Sevilla, D., Gibson, G., Kutt, H., Blau, A., Edelstein, L., Hatch, J., Blass, J., 1995. 3,4-Diaminopyridine as a treatment for amyotrophic lateral sclerosis. *J. Neurol. Sci.* 129, 21–24.

- Aldoss, I.T., Tashi, T., Ganti, A.K., 2009. Seliciclib in malignancies. *Expert Opin. Invest. Drugs* 18, 1957–1965.
- Arnon, S.S., Schechter, R., Inglesby, T.V., Henderson, D.A., Bartlett, J.G., Ascher, M.S., Eitzen, E., Fine, A.D., Hauer, J., Layton, M., Lillibridge, S., Osterholm, M.T., O'Toole, T., Parker, G., Perl, T.M., Russell, P.K., Swerdlow, D.L., Tonat, K., 2001. Botulinum toxin as a biological weapon: medical and public health management. *JAMA* 285, 1059–1070.
- Benson, C., White, J., De Bono, J., O'Donnell, A., Raynaud, F., Cruickshank, C., McGrath, H., Walton, M., Workman, P., Kaye, S., Cassidy, J., Gianella-Borradori, A., Judson, I., Twelves, C., 2007. A phase I trial of the selective oral cyclin-dependent kinase inhibitor seliciclib (CYC202; R-Roscovotine), administered twice daily for 7 days every 21 days. *Br. J. Cancer* 96, 29–37.
- Benson, M.A., Fu, Z., Kim, J.-P., Baldwin, M.R., 2011. Unique ganglioside recognition strategies for clostridial neurotoxins. *J. Biol. Chem.* 286, 34015–34022.
- Bever Jr., C.T., Leslie, J., Camenga, D.L., Panitch, H.S., Johnson, K.P., 1990. Preliminary trial of 3,4-diaminopyridine in patients with multiple sclerosis. *Ann. Neurol.* 27, 421–427.
- Buraei, Z., Angheliescu, M., Elmslie, K.S., 2005. Slowed N-type calcium channel (CaV2.2) deactivation by the cyclin-dependent kinase inhibitor roscovotine. *Biophys. J.* 89, 1681–1691.
- Buraei, Z., Schofield, G., Elmslie, K.S., 2007. Roscovotine differentially affects CaV2 and Kv channels by binding to the open state. *Neuropharmacology* 52, 883–894.
- Cherington, M., Ryan, D.W., 1968. Botulism and guanidine. *N. Engl. J. Med.* 278, 931–933.
- Cherington, M., Schultz, D., 1977. Effect of guanidine, germin and steroids in a case of botulism. *Clin. Toxicol.* 11, 19–25.
- DasGupta, B.R., Sugiyama, H., 1972. A common subunit structure in *Clostridium botulinum* type A, B and E toxins. *Biochem. Biophys. Res. Commun.* 48, 108–112.
- Davis, L.E., Johnson, J.K., Bicknell, J.M., Levy, H., McEvoy, K.M., 1992. Human type A botulism and treatment with 3,4-diaminopyridine. *Electromyogr. Clin. Neurophysiol.* 32, 379–383.
- Delgado-Martínez, I., Nehring, R.B., Sørensen, J.B., 2007. Differential abilities of SNAP-25 homologs to support neuronal function. *J. Neurosci.* 27, 9380–9391.
- DeStefino, N.R., Pilato, A.A., Dittich, M., Cherry, S.V., Cho, S., Stiles, J.R., Meriney, S.D., 2010. (R)-roscovotine prolongs the mean open time of unitary N-type calcium channel currents. *Neuroscience* 167, 838–849.
- Dong, M., Yeh, F., Tepp, W.H., Dean, C., Johnson, E.A., Janz, R., Chapman, E.R., 2006. SV2 is the protein receptor for botulinum neurotoxin A. *Science* 312, 592–596.
- Fischer, A., Montal, M., 2007. Crucial role of the disulfide bridge between botulinum neurotoxin light and heavy chains in protease translocation across membranes. *J. Biol. Chem.* 282, 29604–29611.
- Gu, S., Rumpel, S., Zhou, J., Strotmeier, J., Bigalke, H., Perry, K., Shoemaker, C.B., Rummel, A., Jin, R., 2012. Botulinum neurotoxin is shielded by NTN4 in an interlocked complex. *Science* 335, 977–981.
- Guendel, I., Agbottah, E.T., Kehn-Hall, K., Kashanchi, F., 2010. Inhibition of human immunodeficiency virus type-1 by cdk inhibitors. *Aids Res. Ther.* 7, 1–14.
- Hatheway, C.L., Snyder, J.D., Seals, J.E., Edell, T.A., Lewis Jr., G.E., 1984. Antitoxin levels in botulism patients treated with trivalent equine botulism antitoxin to toxin types A, B, and E. *J. Infect. Dis.* 150, 407–412.
- Hayes, K.C., 2011. Impact of extended-release dalfampridine on walking ability in patients with multiple sclerosis. *Neuropsychiatr. Dis. Treat.* 7, 229–239.
- Howe, J.R., Ritchie, J.M., 1991. On the active form of 4-aminopyridine: block of K^+ currents in rabbit Schwann cells. *J. Physiol. (Lond.)* 433, 183–205.
- Kalia, J., Swartz, K.J., 2011. Elucidating the molecular basis of action of a classic drug: guanidine compounds as inhibitors of voltage gated potassium channels. *Mol. Pharmacol.* 80, 1085–1095.
- Kalla, R., Spiegel, R., Claassen, J., Bardins, S., Hahn, A., Schneider, E., Rettinger, N., Glasauer, S., Brandt, T., Strupp, M., 2011. Comparison of 10-mg doses of 4-aminopyridine and 3,4-diaminopyridine for the treatment of downbeat nystagmus. *J. Neuroophthalmol.* 31, 320–325.
- Korenke, A.R., Rivey, M.P., Allington, D.R., 2008. Sustained-release fampridine for symptomatic treatment of multiple sclerosis. *Ann. Pharmacother.* 42, 1458–1465.
- Korazova, L.K., Montal, M., 2003. Translocation of botulinum neurotoxin light chain protease through the heavy chain channel. *Nat. Struct. Biol.* 10, 13–18.
- Kukreja, R.V., Singh, B.R., 2007. Comparative role of neurotoxin-associated proteins in the structural stability and endopeptidase activity of botulinum neurotoxin complex types A and E. *Biochemistry* 46, 14316–14324.
- Le Tourneau, C., Faivre, S., Laurence, V., Delbaldo, C., Vera, K., Girre, V., Chiao, J., Armour, S., Frame, S., Green, S.R., Gianella-Borradori, A., Diéras, V., Raymond, E., 2010. Phase I evaluation of seliciclib (R-roscovotine), a novel oral cyclin-dependent kinase inhibitor, in patients with advanced malignancies. *Eur. J. Cancer* 46, 3243–3250.
- Li, B., Peet, N.P., Butler, M.M., Burnett, J.C., Moir, D.T., Bowlin, T.L., 2010. Small molecule inhibitors as countermeasures for botulinum intoxication. *Molecules* 16, 202–220.
- Lin, M.J., Lin-Shiau, S.Y., 1997. Multiple types of Ca^{2+} channels in mouse motor nerve terminals. *Eur. J. Neurosci.* 9, 817–823.
- Lindström, M., Korkeala, H., 2006. Laboratory diagnostics of botulism. *Clin. Microbiol. Rev.* 19, 298–314.
- Lundh, H., Leander, S., Thesleff, S., 1977. Antagonism of the paralysis produced by botulinum toxin in the rat. The effects of tetraethylammonium, guanidine and 4-aminopyridine. *J. Neurol. Sci.* 32, 29–43.
- Marcus, S.M., 2009. Reflections on the care of a patient severely poisoned by 'rogue' botulinum toxin and rendered paralysed for a protracted hospital stay. *The Botulinum J.* 1, 318–339.

- Mayorov, A.V., Willis, B., Di Mola, A., Adler, D., Borgia, J., Jackson, O., Wang, J., Luo, Y., Tang, L., Knapp, R.J., Natarajan, C., Goodnough, M.C., Zilberberg, N., Simpson, L.L., Janda, K.D., 2010. Symptomatic relief of botulinum neurotoxin/A intoxication with aminopyridines: a new twist on an old molecule. *ACS Chem. Biol.* 5, 1183–1191.
- Molgó, J., Lemeignan, M., Thesleff, S., 1987. Aminoglycosides and 3,4-diaminopyridine on neuromuscular block cause by botulinum type A toxin. *Muscle Nerve* 10, 464–470.
- Molgó, J., Lundh, H., Thesleff, S., 1980. Potency of 3,4-diaminopyridine and 4-aminopyridine on mammalian neuromuscular transmission and the effect of pH changes. *Eur. J. Pharmacol.* 61, 25–34.
- Montecucco, C., Schiavo, G., Pantano, S., 2005. SNARE complexes and neuroexocytosis: how many, how close? *Trends Biochem.* 30, 367–372.
- Muñoz-Caro, C., Niño, A., 2002. The nature of the receptor site for the reversible K⁺ channel blocking by aminopyridines. *Biophys. Chem.* 96, 1–14.
- Nakamura, Y., Takahashi, T., 2007. Developmental changes in potassium currents at the rat calyx of Held presynaptic terminal. *J. Physiol. (Lond.)* 581, 1101–1112.
- Paton, J.F., 1997. Rhythmic bursting of pre- and post-inspiratory neurons during central apnoea in mature mice. *J. Physiol. (Lond.)* 502, 623–639.
- Penner, R., Dreyer, F., 1986. Two different presynaptic calcium currents in mouse motor nerve terminals. *Pflügers Arch.* 406, 190–196.
- Puggiari, M., Cherington, M., 1978. Botulism and guanidine. Ten years later. *JAMA* 240, 2276–2277.
- Rega, P.P., Bork, C.E., Burkholder-Allen, K., Bisesi, M.S., Gold, J.P., 2010. Single-breath count test: an important adjunct in the triaging of patients in a mass-casualty incident due to botulism. *Prehospital Disaster Med.* 25, 219–222.
- Robinson, R.F., Nahata, M.C., 2003. Management of botulism. *Ann. Pharmacother.* 37, 127–131.
- Ruthel, G., Burnett, J.C., Nuss, J.E., Wanner, L.M., Tressler, L.E., Torres-Melendez, E., Sandwick, S.J., Retterer, C.J., Bavari, S., 2011. Post-intoxication inhibition of botulinum neurotoxin serotype A within neurons by small-molecule, non-peptidic inhibitors. *Toxins* 3, 207–217.
- Shapiro, R.L., Hatheway, C., Swerdlow, D.L., 1998. Botulism in the United States: a clinical and epidemiological review. *Ann. Intern. Med.* 129, 221–228.
- Shi, R., Sun, W., 2011. Potassium blockers as an effective treatment to restore impulse conduction in injured axons. *Neurosci. Bull.* 27, 36–44.
- Simpson, L.L., 2004. Identification of the major steps in botulinum toxin action. *Annu. Rev. Pharmacol. Toxicol.* 44, 167–193.
- Simpson, L.L., 1986. A preclinical evaluation of aminopyridines as putative therapeutic agents in the treatment of botulism. *Infect. Immun.* 52, 858–862.
- Sobel, J., 2005. Botulism. *Clin. Infect. Dis.* 41, 1167–1173.
- Tacket, C.O., Shandera, W.X., Mann, J.M., Hargrett, N.T., Blake, P.A., 1984. Equine antitoxin use and other factors that predict outcome in type A foodborne botulism. *Am. J. Med.* 76, 794–798.
- Thesleff, S., 1989. Botulinum neurotoxins as tools in studies of synaptic mechanisms. *Quart. J. Exp. Physiol.* 74, 1003–1017.
- Tsai, Y.C., Maditz, R., Kuo, C.-L., Fishman, P.S., Shoemaker, C.B., Oyler, G.A., Weissman, A.M., 2010. Targeting botulinum neurotoxin persistence by the ubiquitin-proteasome system. *Proc. Natl. Acad. Sci. USA* 107, 16554–16559.
- Vollmer, T., Blight, A.R., Henney 3rd, H.R., 2009. Steady-state pharmacokinetics and tolerability of orally administered fampridine sustained-release 10-mg tablets in patients with multiple sclerosis: a 2-week, open label, follow-up study. *Clin. Ther.* 31, 2215–2223.
- Yan, Z., Chi, P., Bibb, J.A., Ryan, T.A., Greengard, P., 2002. Roscovitine: a novel regulator of P/Q-type calcium channels and transmitter release in central neurons. *J. Physiol. (Lond.)* 540, 761–770.
- Yarotsky, V., Elmslie, K.S., 2012. Roscovitine inhibits CaV3.1 (T-type) channels by preferentially affecting closed state inactivation. *J. Pharmacol. Exp. Ther.* 340, 463–472.